

Received: 13.02.2012 / Accepted: 06.03.2012

DOI: 10.5137/1019-5149.JTN.6008-12.0

Prevention of Epidural Fibrosis in Rats by Local or Systemic Administration of Citicoline

Sıçanlarda Lokal ve Sistemik Sitikolin Uygulanması ile Epidural Fibrozisin Önlenmesi

Mehmet SAVRAN¹, Ahmet BEKAR¹, Mehmet CANSEV², Sahsene TOLUNAY³, Ismail Hakki ULUS⁴, Mevlut Ozgur TASKAPILIOGLU¹

¹Uludag University, Faculty of Medicine, Department of Neurosurgery, Bursa, Turkey

²Uludag University, Faculty of Medicine, Department of Pharmacology, Bursa, Turkey

³Uludag University, Faculty of Medicine, Department of Pathology, Bursa, Turkey

⁴Acibadem University, Faculty of Medicine, Department of Pharmacology, Istanbul, Turkey

Correspondence address: Ahmet BEKAR / E-mail: abekar@uludag.edu.tr

ABSTRACT

AIM: The objective of this study was to investigate the effect of citicoline administration on epidural fibrosis which is a frequent complication of lumbar disc surgery with no effective treatment or preventive surgical technique.

MATERIAL and METHODS: Sixty Sprague-Dawley female rats undergoing L4-5 right hemilaminotomy and annular fenestration were arranged in three groups: rats in Group 1 (control group) and Group 2 (topical citicoline group) were applied 0,9% saline and 100 µM citicoline on surgical area, respectively, while rats in Group 3 (systemic citicoline group) received 600 µmol/kg citicoline intraperitoneally. Rats were sacrificed four weeks later and their vertebral colons were removed en bloc. Groups were evaluated according to histological criteria and results were compared using statistical tools.

RESULTS: Compared with control group, significantly less epidural fibrosis, dural adhesion, fibroblast cell density, foreign body reaction, and medulla spinalis retraction were observed in groups treated with topical and systemic citicoline (groups 2 and 3) (p<0,001). No significant difference was found with regard to measured parameters between two treatment groups (p>0,05).

CONCLUSION: Our study demonstrates for the first time in the literature that citicoline may be effective for preventing postoperative epidural fibrosis. However, its mechanism of action and clinical effectiveness must be further investigated.

KEYWORDS: Epidural fibrosis, Annular fenestration, Laminotomy, Citicoline, Rat

ÖZ

AMAÇ: Çalışmamızın amacı, sitikolin uygulamasının henüz efektif bir tedavisi veya koruyucu bir cerrahi tekniği bulunmayan ve lomber disk cerrahisinin sık bir komplikasyonu olan epidural fibrozis üzerine etkisini incelemektir.

YÖNTEM ve GEREÇLER: Sağ L4-5 hemilaminotomive anular fenestrasyonu yapılmış 60 Sprague Dawley dişi sıçan 3 gruba ayrıldı. Grup 1 (kontrol grubu) ve Grup 2 (topikal sitikolin grubu)'deki sıçanlara cerrahi alan üzerine sırasıyla 0,9% serum fizyolojik ve 100 µM sitikolin uygulanırken, Grup 3 (sistemik sitikolin grubu)'deki sıçanlara intraperitoneal yolla 600 µmol/kg sitikolin verildi. Sıçanlar 4 hafta sonra öldürüldü ve vertebral kolonları blok olarak çıkarıldı. Gruplar histolojik kriterlere göre değerlendirildi ve sonuçlar istatistiksel yöntemler kullanılarak karşılaştırıldı.

BULGULAR: Topikal ve sistemik sitikolin ile tedavi edilen (Grup 2 ve 3) gruplarda kontrol grubuna kıyasla istatistiksel olarak daha az epidural fibrozis, dural yapışıklık, fibroblast hücre yoğunluğu, yabancı cisim reaksiyonu ve medulla spinalis retraksiyonu gözlemlendi (p<0,001). İncelenen parametreler bakımından iki tedavi grubu arasında anlamlı bir fark bulunmadı (p>0,05).

SONUÇ: Çalışmamız literatürde ilk kez sitikolin'in postoperatif epidural fibrozisi önlemede efektif olabileceğini göstermiştir. Ancak sitikolinin etki mekanizmaları ve klinik etkinliği ilerideki çalışmalarla araştırılmalıdır.

ANAHTAR SÖZCÜKLER: Epidural fibrozis, Annular fenestrasyon, Laminotomi, Sitikolin, Sıçan

INTRODUCTION

Surgical dissection into the spinal canal is challenging during lumbar revision surgery in cases with epidural fibrosis since it increases such intraoperative complications as occurrence of dural tears, spinal nerve lesions, and local bleeding (3, 34, 35, 39, 46). Dense scar tissue increases the morbidity of subsequent operations to the same region (11, 26, 43, 45).

Braverman et al., reported that repeated surgery for fibrosis had only a 30% to 35% success rate (9), whereas 15% to 20% of patients reported worsening of their symptoms (33).

Various surgical procedures including minimally invasive surgery with small incision, limited manipulations during surgery, proper hemostasis and removing foreign materials from epidural region, placing mechanical barriers within dura

and adjacent tissue have been proposed to reduce epidural fibrosis (3, 20, 37). In addition, numerous materials and drugs have been used experimentally and clinically to prevent epidural fibrosis (8, 10, 16, 18, 20, 22, 31, 32, 46, 56). However, epidural fibrosis still remains to be a major clinical problem in patients undergoing surgery for disc herniation.

Cytidine 5'-diphosphate choline (CDP-choline) is an endogenous intermediate in the biosynthesis of cell membrane phospholipids and exogenously-administered CDP-choline is referred to as citicoline (53). Experimental and clinical studies reported beneficial use of citicoline in neurodegenerative diseases such as Alzheimer's and Parkinson's disease, in improving learning and memory, acute and chronic cerebrovascular diseases, traumatic brain injury and glaucoma (30, 47, 48).

Data obtained in our previous studies on citicoline's effects on promoting regeneration of peripheral nerves subjected to immediate section suturing type surgery and reducing postoperative scarring, and recent similar observations encouraged us to test citicoline in prevention of epidural fibrosis after lumbar disc surgery (6, 12, 36). Hence, the purpose of our study was to investigate local and systemic effectiveness of citicoline on epidural fibrosis in a rat model of laminotomy and discectomy.

MATERIAL and METHODS

Rat population and Surgical Technique:

Our study was approved by the Animal Research Ethical Committee of Uludag University. We used 60 female Sprague-Dawley rats weighing 200-250 grams. Each animal received one dose intramuscular (i.m.) injection of cephazolin sodium (20 mg/kg) for prophylaxis. Rats were anesthetized with ketamine hydrochloride (70 mg/kg; i.m.) (Ketalar, Pfizer-Turkey, 2007) plus Xylazine hydrochloride (10 mg/kg; i.m.) (Rompun, Bayer, Turkey). After stabilizing rats on the operation table in prone position, the surgical site was brushed with ChlorHex-OL and sterilized with povidone-iodine solution. Bilateral paraspinal muscles were subperiosteally dissected following a lumbar midline incision. Under operation microscope (Opmi, CARL ZEISS, Germany) the L3-5 laminae were exposed and a right hemilaminectomy were performed at L-4 level using high-speed drill. Dura mater was exposed by removing epidural fat tissue. After finding disc space annular fenestration were performed by using 22G catheter. Hemostasis was maintained by irrigation with saline and surgical gauze throughout the surgical procedure. The hemilaminectomy site was marked with 4-0 vicryl in all rats and the layers were closed. Rats were divided into three groups. Group 1 (control group, n=20) underwent L4-5 right hemilaminotomy and annular fenestration, and spongostan soaked in 0,9% saline was applied to surgical area. In Group 2 (topical citicoline group, n=20) spongostan soaked in citicoline (100 µM) was applied to surgical area locally after identical surgical procedure. In Group 3 (systemic citicoline group, n=20) citicoline was applied intraperitoneally at a dose of 600 µmol/kg after identical surgical procedure.

Preparation of Specimens:

Four weeks after the surgery rats were killed by ether inhalation. For histopathologic studies, the lumbar spine including the surgical site was removed en bloc. The specimen was immersed into 10% buffered formalin solution. Each specimen was then decalcified in equal quantities of 10% formic acid and 8% hydrochloric acid solution for 3 days. All specimens were then washed with tap water for 12 hours. Five ±m thick serial sections were cut from formalin-fixed paraffin-embedded tissues and stained with hematoxylin and eosin (HE) and Masson trichrome (MT).

Histopathological Evaluation:

All specimens were examined microscopically by the same investigator who was blinded to the study. The density of epidural fibrosis, dural adhesion, foreign body reaction, fibroblast cell density and medulla spinalis retraction were investigated. The density of epidural fibrosis, foreign body reaction, and medulla spinalis retraction were scored as absent (0), mild (1), moderate (2), or severe (3) as described previously (49) (Table I).

Dural adhesions were scored as follows: grade 0, the dura is free of scar tissue; grade 1, thin fibrous bands are observed between the dura and scar tissue; grade 2, continuous adherence is observed in less than two-thirds of the laminectomy defect; grade 3, adherence is observed in more than two-thirds of the laminectomy defect, or extends to the nerve roots as described previously (23) (Table II).

Grading of the fibroblast cell density was made as follows: grade 1, less than 100 fibroblast cells per 400x field; grade 2, 100-150 fibroblast cells; grade 3, more than 150 fibroblast cells as described previously (24) (Table III).

Statistical analysis:

Statistical analyses were performed using SPSS 13.0 (Chicago, IL.) software. Median minimum and maximum values were granted for the density of epidural fibrosis, dural adhesion, foreign body reaction, fibroblast cell density and medulla spinalis retraction. Comparisons between groups were made using Kruskal Wallis test. In subgroup analysis, groups were compared in doubles using Mann-Whitney U test and p value <0.05 was considered statistically significant.

RESULTS

Four rats died during the study: two of them due to anesthetic complications and the other two due to dural injury. These 4 animals were replaced in order to achieve a constant number of rats in all three groups.

All rats recovered without wound infection. Similarly, no signs of infection were noted in subcutaneous tissue, fascia, and paravertebral muscles.

Group 1 (control group) showed dense epidural scar formation at laminectomy defects (Figure 1, 2A-C). Compared with group 1, significantly lower values were obtained with

Table I: Scoring of the Density of Epidural Fibrosis, Foreign Body Reaction, and Medulla Spinalis Retraction

Grade 0	Absent
Grade 1	Mild
Grade 2	Moderate
Grade 3	Severe

Table II: Grading of the Extent of Dural Adhesions

Grade 0	The dura is free of scar tissue
Grade 1	Thin fibrous bands are observed between the dura and scar tissue
Grade 2	Continuous adherence is observed in less than two-thirds of the laminectomy defect
Grade 3	The adherence is observed in more than two-thirds of the laminectomy defect, or extends to the nerve roots

Table III: Grading of the Fibroblast Cell Density

Grade 1	Less than 100 (400x)
Grade 2	100 – 150 (400x)
Grade 3	More than 150 (400x)

regard to epidural scar formation ($p < 0,001$ and $p < 0,001$), dural adhesion ($p < 0,001$ and $p < 0,001$), fibroblast cell density ($p < 0,001$ and $p < 0,001$), foreign body reaction ($p = 0,021$ and $p = 0,013$) and medulla spinalis retraction ($p = 0,043$ and $p = 0,033$) in group 2 (topical citicoline group) and group 3 (systemic citicoline group), respectively. On the other hand, considering all parameters measured, no significant differences were found between topical citicoline (group 2) and systemic citicoline (group 3) groups ($p > 0,05$). Grading scores of all parameters in three groups and comparative graphics of mean values are shown in Table IV and Figure 1, respectively.

DISCUSSION

These data show that citicoline treatment significantly reduces epidural scar formation, dural adhesion, fibroblast cell density, foreign body reaction and medulla spinalis retraction in a rat model of laminotomy and discectomy. Similar beneficial effects are observed by either local or systemic administration.

Epidural fibrosis is usually observed post-spinal surgery as nonphysiological scar formation around the origin of the radicular sheath. It behaves as a reparative inflammation causing clinical problems of characteristic nature (5, 32, 51, 52). Epidural fibrosis affects the normal neural dynamics of nerve root by producing adhesions tethering the nerve root to adjacent tissues. Root ischemia caused by tethering may be an important contributor to the generation of radicular pain (44).

Discectomy (annular fenestration) was not performed in most experimental studies that evaluated postoperative scar formation, because of the belief that epidural fibrosis derived from posterior spinal muscles. However, it was reported that nucleus pulposus, which appears in peridural region due to annular fenestration, could take part in the formation of peridural fibrosis and in the etiology of pain by inflammatory immunological effects and by causing microvascular changes (43). Therefore, we used laminotomy and discectomy model in our study to investigate the effect of citicoline on postoperative scarring.

Attempts have been made for prevention of epidural fibrosis after lumbar disc surgery, and, to date, many materials have been used in this regard (3, 8, 22, 29). Despite encouraging results obtained in experimental studies, no mechanical or chemical intervention has proved to be beneficial in the clinical setting so far. Therefore, preventing post operative epidural fibrosis continues to be a clinical problem.

Fat grafts were the first to be tested, and currently they are the most commonly used materials. Unfortunately, these grafts showed little benefit; they reduce tethering rather than preventing epidural fibrosis (22, 40). Numerous materials and strategies including low dose external beam radiation, a carbohydrate polymer, ADCON-L, non-steroidal anti inflammatory drugs (NSAID), anticancer agents such as mitomycin C, cyclosporin A, and 5-fluorouracil, topical thrombolysis with recombinant tissue plasminogen activator (r-tPA), and seprafilm and interceed TC7 have been tested




Figure 1: Comparative graphics of mean values. The values in group 2 (topical citicoline group) and group 3 (systemic citicoline group) were significantly lower than those in group 1 (0,9% saline group). Significance levels between group 1 and 2 with regard to epidural fibrosis, dural adhesion, fibroblast cell density, foreign body reaction and medulla spinalis retraction were $p < 0,001$, $p < 0,001$, $p < 0,001$, $p < 0,021$, and $p < 0,043$, respectively. Significance levels between group 1 and 3 with regard to epidural fibrosis, dural adhesion, fibroblast cell density, foreign body reaction and medulla spinalis retraction were $p < 0,001$, $p < 0,001$, $p < 0,001$, $p < 0,013$, and $p < 0,033$ respectively. There were no significant differences between group 2 and 3 in terms of the measured parameters ($p > 0,05$).




Figure 2: Representative images of Grade 3 (A), Grade 2 (B) and Grade 1 (C) epidural fibrosis. Images on the left panel were obtained by Hematoxylin-eosine (HE) staining and those on the right panel were obtained by Masson's trichrome (MT) staining. **EF:** Epidural fibrosis, **LA:** laminotomy area, **DM:** Dura mater, **MS:** Medulla spinalis.

for preventing epidural fibrosis, however, none of them have entered clinical use (8, 16, 18, 19, 21, 23, 29, 42, 46). Therefore, new studies are required in order to discover new materials or chemicals that could prevent epidural fibrosis.

CDP-choline (cytidine 5'- diphosphocholine), is an endogenous intermediate in the biosynthesis of cell membrane phospholipids and exogenously-administered CDP-choline is referred to as citicoline (53). Citicoline and its metabolite choline serve as substrates for the synthesis of phosphatidylcholine, a primary component of neuronal membrane (2, 17, 53). Grouped with the B vitamins, choline is a trimethylated nitrogenous base that enters three major metabolic pathways: (1) phospholipid synthesis via phosphorylcholine; (2) acetylcholine synthesis; and (3) oxidation to betaine, which serves as a methyl donor (17).

Citicoline has been shown to exhibit beneficial effects in several neurological conditions. For example, citicoline improves neurological outcomes and reduces the volume of ischemic injury in patients with stroke (17, 41). In addition, citicoline reduces infarct size and decreases mortality in

hypotensive rats with subarachnoid hemorrhage and basilar artery occlusion (4).

The observed pathologies with regard to epidural fibrosis in the present rat laminotomy and discectomy model suggest the onset of inflammatory processes after the surgery. Hence, treatments targeting the inflammatory response might confer benefit in lumbar disc surgery. We found reduced scar formation accompanied by decreased dural adhesion, fibroblast cell density, foreign body reaction and medulla spinalis retraction following either local or systemic administration of citicoline in our study. The mechanism(s) whereby citicoline exhibits such effects are not yet known. However, evidence for possible citicoline mechanism of action has been provided in previous studies. For example, citicoline was reported to decrease phospholipase A2 stimulation thereby leading to blockade of generation of inflammatory prostaglandins and thromboxanes (1).

Although these anti-inflammatory effects seem to have resulted from direct action of citicoline, it is also possible that choline metabolized from citicoline might have contributed

Table IV: Grading Scores of all Parameters in Three Groups

Rat No	Density of Epidural fib.			Dural Adhesion			Fibroblast Cell Density			Foreign Body Reaction			Medulla Spinalis Retraction		
	Group 1	Group 2	Group 3	Group 1	Group 2	Group 3	Group 1	Group 2	Group 3	Group 1	Group 2	Group 3	Group 1	Group 2	Group 3
1	3	2	2	3	2	2	3	2	2	3	2	0	0	0	3
2	3	2	1	3	2	1	3	1	1	3	0	0	3	1	0
3	3	2	2	2	2	2	3	1	1	2	0	0	1	0	0
4	3	1	1	2	1	2	3	1	1	3	0	0	0	0	0
5	2	1	1	2	1	1	3	1	1	1	1	0	2	0	0
6	3	2	1	3	2	1	3	2	1	0	0	0	2	0	0
7	3	2	1	3	2	1	3	2	1	0	1	1	0	0	1
8	3	2	1	3	1	1	2	3	2	1	0	0	0	0	0
9	3	1	2	3	2	1	1	2	1	1	1	0	0	0	0
10	3	2	1	3	2	1	3	2	1	2	0	0	1	0	0
11	3	2	2	3	2	1	2	2	1	0	0	0	2	0	0
12	3	1	1	3	1	2	3	1	1	2	0	1	1	2	0
13	3	1	2	3	1	2	2	1	1	2	0	1	1	0	0
14	3	1	1	3	2	1	3	2	1	2	1	1	0	0	0
15	2	1	2	3	1	1	3	2	1	1	2	1	0	0	0
16	3	2	1	2	1	2	1	1	2	2	0	1	0	0	0
17	3	1	1	3	2	2	3	1	1	1	0	1	0	1	0
18	3	2	1	3	1	1	3	2	1	0	1	1	2	0	0
19	2	1	2	3	2	1	2	1	1	0	0	0	3	0	0
20	2	1	1	2	1	1	1	1	1	0	1	1	0	0	0

to the observed therapeutic effects. Recent studies demonstrated anti-inflammatory properties of cholinergic activation and the term "cholinergic antiinflammatory pathway" has been proposed (50). CDP-choline is a choline donor since part of exogenous CDP-choline is metabolized to choline (14, 54). While choline deficiency results in increased levels of proinflammatory cytokines such as TNF- α (25), parenteral choline administration attenuates endotoxin-induced multiple organ injury and the elevation in circulating TNF- α levels (27). Since CDP-choline is a source of choline and its administration leads to initiation of responses in cholinergic nature (15, 28), it could be speculated that one mechanism by which CDP-choline ameliorates the severity of scar formation after lumbar disc surgery might be the action of choline that was released by citicoline breakdown on inflammatory processes. This hypothesis is further supported by evidence from a previous study in which we demonstrated beneficial effects of both citicoline and choline on reducing postoperative scarring in a rat model of peripheral nerve injury (6). In addition, reversal of mechanical hyperalgesia in carrageenan-induced inflammatory pain model after citicoline administration (7) was mediated in part by supraspinal $\alpha 7$ -selective nicotinic ACh receptors which utilizes choline as a selective agonist (38). Therefore, our present

observations could have resulted from possible blockade of the inflammatory phase of scar formation by citicoline or its metabolite choline. These possible mechanisms are now being tested in our laboratory.

CONCLUSIONS

To the best of our knowledge, this is the first experimental study demonstrating prevention of epidural fibrosis by citicoline in a rat model of laminotomy and discectomy. Our present data show that citicoline effectively prevents epidural fibrosis by either local or systemic administration. Future studies are warranted in order to determine citicoline's mechanisms of action on epidural fibrosis and to test its clinical efficiency.

REFERENCES

1. Adibhatla RM, Hatcher JF: Citicoline decreases phospholipase A2 stimulation and hydroxyl radical generation in transient cerebral ischemia. *J Neurosci Res* 73:308-315, 2003
2. Agut J, Lopez G.-Coviella I, Ortiz JA, Wurtman RJ: Oral cytidine 5'-diphosphate choline administration to rats increases brain phospholipid levels. *Ann NY Acad Sci* 695:318-320, 1993
3. Alkalay RN, Kim DH, Urry DW, Xu J, Parker TM, Glazer PA: Prevention of postlaminectomy epidural fibrosis using bioelastic materials. *Spine* 28:1659-1665, 2003

4. Alkan T, Kahveci N, Goren B, Korfali E, Ozluk K: Ischemic brain injury caused by interrupted versus uninterrupted occlusion in hypotensive rats with subarachnoid hemorrhage: Neuroprotective effects of citicoline. *Arch Physiol Biochem* 109:161-167, 2001
5. Asch HL, Lewis PJ, Moreland DB, Egnatchik JG, Yu YJ, Clabeaux DE, Hyland AH: Prospective multiple outcomes study of outpatient lumbar microdiscectomy: Should 75-80% success rates be the norm? *J Neurosurg* 96:34-44, 2002
6. Aslan E, Kocaeli H, Bekar A, Tolunay S, Ulus IH: CDP-choline and its endogenous metabolites, cytidine and choline, promote the nerve regeneration and improve the functional recovery of injured rat sciatic nerves. *Neurol Res* 33:766-773, 2011
7. Bagdas D, Sonat FA, Hamurtekin E, Sonal S, Gurun MS: The antihyperalgesic effect of cytidine-5'-diphosphate-choline in neuropathic and inflammatory pain models. *Behav Pharmacol* 22:589-598, 2011
8. Boyaci S, Bekar A, Kocaeli H, Doygun M, Tolunay S: The effectiveness of Adcon-L in preventing peridural fibrosis after the spinal surgery. *Turk Norosirurji Dergisi* 10:102-108, 2000
9. Braverman DL, Slipman CW, Lendrow DA: Using gabapentin to treat failed back surgery syndrome caused by epidural fibrosis: A report of two cases. *Arch Phys Med Rehabil* 82: 691-693, 2001
10. Bulut G, Dogan S, Doygun M: The Effectiveness of omental graft in preventing epidural fibrosis after the spinal surgery: An experimental study. *Uludag Universitesi Tip Fakultesi Dergisi* 3:115-121, 2007
11. Burton CV, Kirkaldy-Willis WH, Yong-Hing K, Heithoff KB: Causes of failure of surgery on the lumbar spine. *Clin Orthop Relat Res* 157:191-199, 1981
12. Caner B, Kafa IM, Bekar A, Kurt A, Karli N, Cansev M, Ulus IH: Intraperitoneal administration of CDP-choline or a combination of cytidine plus choline improves nerve regeneration and functional recovery in a rat model of sciatic nerve injury. *Neurol Res (In Press)*
13. Cansev M: Uridine and cytidine in the brain: Their transport and utilization. *Brain Res Rev* 52: 389-397, 2006
14. Cansev M, Yilmaz MS, Ilcol YO, Hamurtekin E, Ulus IH: Cardiovascular effects of CDP-choline and its metabolites: Involvement of peripheral autonomic nervous system. *Eur J Pharmacol* 577:129-142, 2007
15. Cansev M, Ilcol YO, Yilmaz MS, Hamurtekin E, Ulus IH: Choline, CDP-choline or phosphocholine increases plasma glucagon in rats: Involvement of the peripheral autonomic nervous system. *Eur J Pharmacol* 589:315-322, 2008
16. Cemil B, Tun K, Kaptanoglu E, Kaymaz F, Cevirgen B, Comert A, Tekdemir I: Use of pimecrolimus to prevent epidural fibrosis in a postlaminectomy rat model. *Neurosurg Spine* 11: 758-763, 2009
17. Conant R, Schauss AG: Therapeutic applications of citicoline for stroke and cognitive dysfunction in the elderly: A review of the literature. *Altern Med Rev* 9:17-31, 2004
18. Dogan S, Taskapilioglu O, Sahin S, Korfali E: The effects of seprafilm and interceed TC7 on epidural fibrosis in a rat hemilaminectomy model. *Neurosurgery Q* 19:190-195, 2009
19. Einhaus SL, Robertson JT, Dohan FC Jr, Wujek JR, Ahmad S: Reduction of peridural fibrosis after lumbar laminotomy and discectomy in dogs by a resorbable gel (ADCON-L). *Spine* 22:1440-1446, 1997
20. Fritsch EW, Heisel J, Rupp S: The failed back surgery syndrome. Reasons, intraoperative findings and long term results. A report of 182 operative treatments. *Spine* 21:626-633, 1996
21. Gerszten PC, Moossy, JJ, Flickinger JC, Gerszten K, Kalend A, Martinez AJ: Inhibition of peridural fibrosis after laminectomy using low-dose external beam radiation in a dog model. *Neurosurgery* 46:1478-1485, 2000
22. Gorgulu A, Simsek O, Cobanoglu S, Imer M, Parsak T: The effect of epidural free fat graft on the outcome of lumbar disc surgery. *Neurosurg Rev* 27:181-184, 2004
23. He Y, Revel M, Loty B: A quantitative model of post-laminectomy scar formation. Effects of a nonsteroidal anti-inflammatory drug. *Spine* 20:557-563, 1995
24. Hinton JL Jr, Warejcka DJ, Mei Y, McLendon RE, Laurencin C, Lucas PA, Robinson JS Jr: Inhibition of epidural scar formation after lumbar laminectomy in the rat. *Spine* 20:564-570, 1995
25. Holmes-McNary MQ, Baldwin AS Jr, Zeisel SH: Opposing regulation of choline deficiency-induced apoptosis by p53 and nuclear factor κ B. *J Biol Chem* 276:41197-41204, 2001
26. Hoyland JA, Freemont AJ, Denton J, Thomas AMC, McMillan JJ, Jayson MIV: Retained surgical swab debris in post-laminectomy arachnoiditis and peridural fibrosis. *J Bone Joint Surg Br* 70:659-662, 1988
27. Ilcol YO, Yilmaz Z, Ulus IH: Endotoxin alters serum-free choline and phospholipid-bound choline concentrations, and choline administration attenuates endotoxin-induced organ injury in dogs. *Shock* 24:288-293, 2005
28. Ilcol YO, Cansev M, Yilmaz MS, Hamurtekin E, Ulus IH: Peripheral administration of CDP-choline and its cholinergic metabolites increases serum insulin: Muscarinic and nicotinic acetylcholine receptors are both involved in their actions. *Neurosci Lett* 431:71-76, 2008
29. Kemaloglu S, Ozkan U, Yilmaz F, Nas K, Gur A, Acemoglu H, Karasu H, Cakmak E: Prevention of spinal epidural fibrosis by recombinant tissue plasminogen activator in rats. *Spinal Cord* 41:427-431, 2003
30. Krupinski J, Ferrer I, Barrachina M, Secades JJ, Mercadal J, Lozano R: CDP-choline reduces pro-caspase and cleaved caspase-3 expression, nuclear DNA fragmentation, and specific PARP-cleaved products of caspase activation following middle cerebral artery occlusion in the rat. *Neuropharmacology*, 42: 846-854, 2002
31. Liu S, Boutrand JP, Tadie M: Use of collagen-based sealant to prevent in vivo epidural adhesions in an adult rat laminectomy model. *J Neurosurg* 94: 61-67, 2001
32. Lladó A, Guimerá J, Garcia F, Navarro A: Expanded polytetrafluoroethylene membrane for the prevention of peridural fibrosis after spinal surgery: An experimental study. *Eur Spine J* 8:138-143, 1999

33. Loupasis GA, Stamos K, Katonis PG, Sapkas G, Korres DS, Hartofilakidis G: Seven- to 20 – year outcome of lumbar discectomy. *Spine* 24:2313-2317, 1999
34. Massie JB, Huang B, Malkmus S, Yaksh TL, Kim CW, Garfin SR, Akeson WH: A preclinical post laminectomy rat model mimics the human post laminectomy syndrome. *J Neurosci Methods* 137:283-289, 2004
35. North RB, Campbell JN, Jams CS, Conover-Walker MK, Wang H, Piantadosi S, Rybock JD, Long DM: Failed back surgery syndrome: 5-year follow-up in 102 patients undergoing repeated operation. *Neurosurgery* 28:685-691, 1991
36. Ozay R, Bekar A, Kocaeli H, Karli N, Filiz G, Ulus IH: Citicoline improves functional recovery, promotes nerve regeneration, and reduces postoperative scarring after peripheral nerve surgery in rats. *Surg Neurol* 68:615-622, 2007
37. Ozer AF, Oktenoglu T, Sasani M, Bozkus H, Canbulat N, Karaarslan E, Sungurlu SF, Sarioglu AC: Preserving the ligamentum flavum in lumbar discectomy: A new technique that prevents scar tissue formation in the first 6 months postsurgery. *Neurosurgery* 59:126-133, 2006
38. Papke RL, Bencherif M, Lippiello P: An evaluation of neuronal nicotinic acetylcholine receptor activation by quaternary nitrogen compounds indicates that choline is selective for the alpha 7 subtype. *Neurosci Lett* 213:201-204, 1996
39. Pospiech J, Pajonk F, Stolke D: Epidural scar tissue formation after spinal surgery: An experimental study. *Eur Spine* 4: 213-219, 1995
40. Prusick VR, Lint DS, Bruder J: Cauda equina syndrome as a complication of free epidural fat-grafting. A report of two cases and a review of the literature. *J Bone Joint Surg Am* 70:1256-1258, 1988
41. Rao AM, Hatcher JF, Dempsey RJ: Lipid alterations in transient forebrain ischemia: Possible new mechanisms of CDP-choline neuroprotection. *J Neurochem* 75:2528-2535, 2000
42. Richter HP, Kast E, Tomczak R, Besenfelder W, Gaus W: Results of applying ADCON-L gel after lumbar discectomy: The German ADCON-L study. *J Neurosurg* 95:179-189, 2001
43. Robertson JT: Role of peridural fibrosis in the failed back: A review. *Eur Spine* J 5:2-6, 1996
44. Rodgers KE, Robertson JT, Espinoza T, Oppelt W, Cortese S, diZerega GS, Berg RA: Reduction of epidural fibrosis in lumbar surgery with Oxiplex adhesion barriers of carboxymethylcellulose and polyethylene oxide. *Spine* J 3:277-283, 2003
45. Ross JS, Zepp R, Modic MT: The postoperative lumbar spine: Enhanced MR evaluation of the intervertebral disk. *AJNR Am J Neuroradiol* 17:323-331, 1996
46. Sandoval MA, Hernandez-Vaquero D: Preventing peridural fibrosis with nonsteroidal anti-inflammatory drugs. *Eur Spine* J 17:451-455, 2008
47. Secades JJ: CDP-Choline: Update and review of its pharmacology and clinical use. *Methods Find Exp Clin Pharmacol* 24:1-53, 2002
48. Sobrado M, Lopez MG, Carceller F, Garcia AG, Roda JM: Combined nimodipine and citicoline reduce infarct size, attenuate apoptosis and increase bcl-2 expression after focal cerebral ischemia. *Neuroscience*, 118: 107-113, 2003
49. Topsakal C, Akpolat N, Erol FS, Ozveren MF, Akdemir I, Kaplan M, Tiftikci M, Kilic N: Sefrafilm superior to Gore-Tex in the prevention of peridural fibrosis. *J Neurosurg* 101:295-302, 2004
50. Tracey KJ: Physiology and immunology of the cholinergic antiinflammatory pathway. *J Clin Invest* 117:289-296, 2007
51. Vogelsang JP, Finkenstaedt M, Vogelsang M, Markakis E: Recurrent pain after lumbar discectomy: The diagnostic value of peridural scar on MRI. *Eur Spine* J 8:475-479, 1999
52. Waguespack A, Schofferman J, Slosar P, Reynolds J: Etiology of long-term failures of lumbar spine surgery. *Pain Med* 3: 18-22, 2002
53. Weiss GB: Minireview. Metabolism and actions of CDP-choline as an endogenous compound and administered exogenously as citicoline. *Life Sci* 56:637-660, 1995
54. Wurtman RJ, Regan M, Ulus I, Yu L: Effect of oral CDP-choline on plasma choline and uridine levels in humans. *Biochem Pharmacol* 60:989-992, 2000
55. Wurtman RJ, Cansev M, Ulus IH: Choline and its products acetylcholine and phosphatidylcholine. In: Lajtha A, ed. *Handbook of Neurochemistry and Molecular Neurobiology: Neural Lipids*. Berlin Heidelberg: Springer-Verlag, 2009: 443-501
56. Yildiz KH, Gezen F, Is M, Cukur S, Dosoglu M: Mitomycin C, 5-fluorouracil, and cyclosporin A prevent epidural fibrosis in an experimental laminectomy model. *Eur Spine* J 16: 1525-1530, 2007